Noname manuscript No.
(will be inserted by the editor)

A nonlinear Korn inequality based on the Green-Saint Venant strain tensor

Alessandro Musesti

arXiv:1605.06964v1 [math-ph] 23 May 2016

May 24, 2016

Abstract A nonlinear Korn inequality based on the Green-Saint Venant strain tensor is proved, whenever the displacement is in the Sobolev space $W^{1,p}$, $p \geq 2$, under Dirichlet conditions on a part of the boundary. The inequality can be useful in proving the coercivity of a nonlinear elastic energy.

Keywords nonlinear Korn inequality, geometric rigidity lemma, finite elasticity, coercivity

Mathematics Subject Classification (2010) 74B20 · 74A05

1 Introduction

Korn inequality is one of the pillars of linear elasticity and it is well-known for more than a century. In the most classical version it writes

$$\|\nabla \boldsymbol{u}\|_2 \leq c\|e(\boldsymbol{u})\|_2 \tag{1}$$

provided that the displacement $\boldsymbol{u}$ vanishes on a sufficiently large part of the boundary. Here $e(\boldsymbol{u})$ denotes the symmetric part of $\nabla \boldsymbol{u}$. Since $e(\boldsymbol{u})$ is the measure of the strain in linear theories, Korn inequality provides a control of the deformation gradient by means of the strain. The L^p version of the inequality, namely

$$\|\nabla \boldsymbol{u}\|_p \leq c\|e(\boldsymbol{u})\|_p \tag{2}$$

if $\boldsymbol{u}$ vanishes on a part of the boundary, followed more recently by general results about singular integral operators, see [6].

However, in nonlinear elasticity a typical measure of the strain is given by the so-called Green-Saint Venant (or Green-Lagrange) strain tensor

$$\mathsf{E}(\boldsymbol{u}) = \frac{1}{2}(\mathsf{F}^{\mathsf{T}}\mathsf{F} - \mathsf{I}) = \frac{1}{2}(\nabla \boldsymbol{u} + \nabla \boldsymbol{u}^{\mathsf{T}} + \nabla \boldsymbol{u}^{\mathsf{T}} \nabla \boldsymbol{u}),$$

A. Musesti
Dipartimento di Matematica e Fisica "Niccolò Tartaglia", Università Cattolica del Sacro Cuore, Via dei Musei 41, I-25121 Brescia (Italy)
E-mail: alessandro.musesti@unicatt.it

where $\mathsf{F} = \nabla \boldsymbol{u}$ is the deformation gradient and I denotes the identity tensor (see for instance [1, Sect. 1.8]). Hence for a hyperelastic material it is customary to write the energy density as a function of $\mathsf{E}(\boldsymbol{u})$. Unfortunately, such an assumption, although very reasonable from a mechanical viewpoint, is a source of difficulties on the mathematical side: both the weakly lower semicontinuity and the coercivity of the elastic energy can be a hard task, or even fail to hold.

Specifically, in proving an existence result for a hyperelastic body subject to external loads and mixed boundary conditions, the Direct Method of the Calculus of Variations can be a very powerful tool. Therefore, even if the elastic energy density is well-behaved (for instance if it is smooth and polyconvex), it is crucial to have some coercivity conditions in order to construct a *bounded* minimizing sequence. Hence it is very important to control the norm of the displacement by the energy. Ultimately, one needs to control the displacement by the Green-Saint Venant tensor, that is, a nonlinear version of the Korn inequality with boundary conditions. The inequality can be useful also in other applications: see for instance [7], where a similar tool allows to prove the existence of a minimizer for a nonlinear elastic problem by the implicit function theorem.

In the present note we prove the inequality

$$\|\nabla \boldsymbol{u}\|_p^p \leq c \|\mathsf{E}(\boldsymbol{u})\|_{p/2}^{p/2} \tag{3}$$

in the case of a bounded Lipschitz body and for $\boldsymbol{u}$ in the Sobolev space $W^{1,p}$, $p \geq 2$, and $\det(\nabla \boldsymbol{u}(\boldsymbol{x}) + \mathsf{I}) > 0$. The result holds under weak assumptions on the boundary conditions, namely it is enough for the displacement $\boldsymbol{u}$ to vanish on a part of the boundary with positive surface measure. We notice that, with a perhaps more simple notation, equation (3) can be rewritten as

$$\|\mathsf{F} - \mathsf{I}\|_p^p \leq C \|\mathsf{F}^{\mathsf{T}}\mathsf{F} - \mathsf{I}\|_{p/2}^{p/2}, \qquad p \geq 2, \ \det \mathsf{F} > 0.$$

Interestingly, as observed by the anonymous reviewer, (2) does not follow from (3) by linearization, since the exponent on the right-hand side is $p/2$.

A similar inequality has been proved in [2] for the case $p = 2$. Here we generalize that result to the case $p \geq 2$. An essential ingredient of the proof is the L^p version of the celebrated *geometric rigidity lemma* of [5], originally stated for L^2, which has been generalized for $1 < p < \infty$ in Conti & Schweizer [3, Sec. 2.4].

2 Notation

Given an $n \times n$-matrix F, we will denote by $|\mathsf{F}|$ its Frobenius norm, that is

$$|\mathsf{F}| = \sqrt{\operatorname{tr}(\mathsf{F}^{\mathsf{T}}\mathsf{F})} = \sum_{i,j=1}^{n} F_{ij}^2,$$

where F^{T} denotes the transpose of the matrix F. We will denote by $SO(n)$ the set of rotations in $\mathbb{R}^n$, that is

$$\mathsf{R} \in SO(n) \quad \Leftrightarrow \quad \mathsf{R}^{\mathsf{T}}\mathsf{R} = \mathsf{I} \text{ and } \det \mathsf{R} = 1,$$

Let Ω be a bounded connected open subset of $\mathbb{R}^n$, $n \geq 2$, with Lipschitz boundary. Concerning the displacement, we assume that $\boldsymbol{u} \in W^{1,p}(\Omega;\mathbb{R}^n)$ with $2 \leq p < \infty$, endowed with the usual norm

$$\|\boldsymbol{u}\|_{1,p} = \left(\|\boldsymbol{u}\|_p^p + \|\nabla \boldsymbol{u}\|_p^p\right)^{1/p}.$$

We will consider homogeneous Dirichlet boundary conditions on a part of the boundary: let $\Gamma \subset \partial\Omega$ be a subset of the boundary of Ω such that

$$|\Gamma| := \mathcal{H}^{n-1}(\Gamma) > 0,$$

where $\mathcal{H}^{n-1}$ denotes the $(n-1)$-dimensional Hausdorff measure. We introduce the set

$$W^{1,p}_{\Gamma} := \left\{ \boldsymbol{u} \in W^{1,p}(\Omega;\mathbb{R}^n) : \ \boldsymbol{u}(\boldsymbol{x}) = \boldsymbol{0} \quad \mathcal{H}^{n-1}\text{-a.e. in } \Gamma \right\}$$

endowed with the norm $\|\cdot\|_{1,p}$, where the equality has to be understood in the sense of traces of Sobolev functions. Notice that $W^{1,p}_{\Gamma}$ is a closed linear subspace of $W^{1,p}(\Omega;\mathbb{R}^n)$.

A crucial lemma is the following (see also [2, Lemma 1]).

Lemma 1 *There is a constant $c > 0$ such that*

$$\operatorname{dist}(\mathsf{F}, SO(n)) \leq c|\mathsf{F}^{\mathsf{T}}\mathsf{F} - \mathsf{I}|^{1/2} \tag{4}$$

for any matrix F with $\det \mathsf{F} > 0$.

Proof Since $\det \mathsf{F} > 0$, it is a well-known consequence of the Polar Factorization Theorem (see [1, Theorem 3.2-2]) that

$$\operatorname{dist}(\mathsf{F}, SO(n)) = |(\mathsf{F}^{\mathsf{T}}\mathsf{F})^{1/2} - \mathsf{I}|.$$

Moreover, denoting by $v_1, \ldots, v_n$ the eigenvalues of the matrix $(\mathsf{F}^{\mathsf{T}}\mathsf{F})^{1/2}$, by the equivalence of the Frobenius and the spectral norm it follows that

$$c_1 \max_{1\leq i\leq n} |v_i - 1| \leq |(\mathsf{F}^{\mathsf{T}}\mathsf{F})^{1/2} - \mathsf{I}| \leq c_2 \max_{1\leq i\leq n} |v_i - 1|$$

for some $c_1, c_2 > 0$. Since $v_1, \ldots, v_n \geq 0$, one has

$$|(\mathsf{F}^{\mathsf{T}}\mathsf{F})^{1/2} - \mathsf{I}| \leq c_2 \max_{1\leq i\leq n} |v_i - 1| \leq c_2 \max_{1\leq i\leq n} |v_i^2 - 1|^{1/2} = \frac{c_2}{c_1}|\mathsf{F}^{\mathsf{T}}\mathsf{F} - \mathsf{I}|^{1/2}$$

and the proof is complete.

3 The nonlinear Korn inequality

We begin by proving the following lemma, which can also be found in [4, Lemma 3.3].

Lemma 2 *Let $\overline{K}$ denote the closure of the cone*

$$K = \Big\{ t(\mathsf{I} - \mathsf{R}) : \ t > 0, \ \mathsf{R} \in SO(n) \Big\}.$$

Then there exists $c > 0$ such that

$$\forall \mathsf{F} \in \overline{K} : \quad |\mathsf{F}| \leq c \min_{\boldsymbol{z} \in \mathbb{R}^n} \left(\int_\Gamma |\mathsf{F}\boldsymbol{x} - \boldsymbol{z}|^2 \, d\mathcal{H}^{n-1}(\boldsymbol{x}) \right)^{1/2}.$$

Proof First of all we notice that

$$\overline{K} = K \cup \mathrm{Skw}(n), \tag{5}$$

where $\mathrm{Skw}(n)$ denotes the set of skew-symmetric tensors of order n. Indeed, assume that $\mathsf{F} \in \overline{K} \setminus K$ with $\mathsf{F} \neq \mathsf{0}$; then there is an unbounded sequence (t_h) with $t_h > 0$ such that

$$t_h(\mathsf{I} - \mathsf{R}_h) \to \mathsf{F} \quad \text{as } h \to \infty,$$

where $\mathsf{R}_h \in SO(n)$. In particular, $|\mathsf{I} - \mathsf{R}_h| \sim 1/t_h$ as $h \to \infty$. Representing an orthogonal tensor as the exponential of a skew-symmetric tensor, we can write

$$\mathsf{R}_h = \exp\left(t_h^{-1}\mathsf{A}\right) + o(t_h^{-1}) \quad \text{as } h \to \infty,$$

for some $\mathsf{A} \in \mathrm{Skw}(n)$. Then we have

$$\mathsf{F} = \lim_{h\to\infty} t_h \left[\mathsf{I} - \exp\left(t_h^{-1}\mathsf{A}\right) - o(t_h^{-1}) \right] = -\frac{d}{dt} \exp(t\mathsf{A}) \Big|_{t=0} = -\mathsf{A},$$

hence F is skew-symmetric.

By (5) it follows that

$$\mathsf{F} \in \overline{K}, \, \mathsf{F} \neq 0 \quad \Rightarrow \quad \dim\ker \mathsf{F} \leq n-2.$$

Moreover it is easy to check that

$$\int_\Gamma |\mathsf{F}\boldsymbol{x} - \boldsymbol{z}|^2 \, d\mathcal{H}^{n-1}(\boldsymbol{x}) \quad \text{has minimum for} \quad \overline{\boldsymbol{z}} = \frac{1}{|\Gamma|} \int_\Gamma \mathsf{F}\boldsymbol{x} \, d\mathcal{H}^{n-1}(\boldsymbol{x}).$$

Now, assume by contradiction that for any $j \geq 1$ there exists $\mathsf{F}_j \in \overline{K}$ such that $|\mathsf{F}_j| = 1$ and

$$\frac{1}{j} > \int_\Gamma |\mathsf{F}_j\boldsymbol{x} - \overline{\boldsymbol{z}}_j|^2 \, d\mathcal{H}^{n-1}(\boldsymbol{x}), \quad \overline{\boldsymbol{z}}_j = \frac{1}{|\Gamma|} \int_\Gamma \mathsf{F}_j\boldsymbol{x} \, d\mathcal{H}^{n-1}(\boldsymbol{x}).$$

Then there exists $\mathsf{F} \in \overline{K}$ such that $|\mathsf{F}| = 1$ and $\mathsf{F}_j \to \mathsf{F}$ up to a subsequence. By continuity it follows that

$$\int_\Gamma |\mathsf{F}\boldsymbol{x} - \overline{\boldsymbol{z}}|^2 \, d\mathcal{H}^{n-1}(\boldsymbol{x}) = 0, \quad \overline{\boldsymbol{z}} = \frac{1}{|\Gamma|} \int_\Gamma \mathsf{F}\boldsymbol{x} \, d\mathcal{H}^{n-1}(\boldsymbol{x}),$$

whence $\mathsf{F}\boldsymbol{x} = \overline{\boldsymbol{z}}$ for $\mathcal{H}^{n-1}$-a.e. $\boldsymbol{x} \in \Gamma$.

Since Γ has positive measure and Ω is Lipschitz, then there exist at least $n-1$ pairs $(\boldsymbol{x}_i, \boldsymbol{y}_i)$ such that $\boldsymbol{x}_i, \boldsymbol{y}_i \in \Gamma$ and $(\boldsymbol{x}_i - \boldsymbol{y}_i)$ are linearly independent. In particular, $\mathsf{F}(\boldsymbol{x}_i - \boldsymbol{y}_i) = 0$, hence $\dim\ker \mathsf{F} \geq n-1$, a contradiction.

Now we are ready to state and prove the main theorem.

Theorem 1 *Let Ω be a bounded connected open subset of $\mathbb{R}^n$ with Lipschitz boundary and $2 \le p < \infty$. Let $\Gamma \subset \partial\Omega$ be a subset of the boundary of Ω such that $\mathcal{H}^{n-1}(\Gamma) > 0$.*

Then there exists a constant $c > 0$ such that

$$\|\nabla \boldsymbol{u}\|_p^p \le c \|\mathsf{E}(\boldsymbol{u})\|_{p/2}^{p/2}$$

for every $\boldsymbol{u} \in W^{1,p}_\Gamma$ such that $\det(\nabla \boldsymbol{u}(\boldsymbol{x}) + \mathsf{I}) > 0$ for a.e. $\boldsymbol{x} \in \Omega$.

Proof The continuity of the trace operator gives

$$\int_\Gamma |f(\boldsymbol{x})|^p \, d\mathcal{H}^{n-1}(\boldsymbol{x}) \le c_1 \int_\Omega (|f(\boldsymbol{x})|^p + |\nabla f|^p) \, d\boldsymbol{x} \tag{6}$$

for every $f \in W^{1,p}(\Omega)$, where $c_1 > 0$ is some constant depending on Ω and Γ.

Let $\boldsymbol{u} \in W^{1,p}_\Gamma$ and consider the vector field $\boldsymbol{\phi} \in W^{1,p}(\Omega; \mathbb{R}^n)$ defined as $\boldsymbol{\phi}(\boldsymbol{x}) = \boldsymbol{u}(\boldsymbol{x}) + \boldsymbol{x}$. For every $\mathsf{R} \in SO(n)$ define

$$\overline{\boldsymbol{z}}_{\mathsf{R}} := \frac{1}{|\Omega|} \int_\Omega (\boldsymbol{\phi}(\boldsymbol{x}) - \mathsf{R}\boldsymbol{x}) \, d\boldsymbol{x}.$$

By (6) and Poincaré-Wirtinger inequality one has

$$\int_\Gamma |\boldsymbol{\phi}(\boldsymbol{x}) - \mathsf{R}\boldsymbol{x} - \overline{\boldsymbol{z}}_{\mathsf{R}}|^p \, d\mathcal{H}^{n-1}(\boldsymbol{x}) \le c_2 \int_\Omega |\nabla \boldsymbol{\phi}(\boldsymbol{x}) - \mathsf{R}|^p \, d\boldsymbol{x}. \tag{7}$$

Notice that in the left-hand side one can replace $\boldsymbol{\phi}(\boldsymbol{x})$ with the identity, in view of the boundary conditions. Moreover, by Lemma 2 it follows that

$$|\mathsf{I} - \mathsf{R}| \le c_3 \left(\min_{\boldsymbol{z} \in \mathbb{R}^n} \int_\Gamma |(\mathsf{I} - \mathsf{R})\boldsymbol{x} - \boldsymbol{z}|^2 \, d\mathcal{H}^{n-1}(\boldsymbol{x}) \right)^{1/2}. \tag{8}$$

In particular, choosing $\boldsymbol{z} = \overline{\boldsymbol{z}}_{\mathsf{R}}$, by Hölder inequality (with $p \ge 2$) and (7) it follows that

$$|\mathsf{I} - \mathsf{R}|^p \le c_3^p |\Gamma|^{\frac{p}{2}-1} \int_\Gamma |(\mathsf{I} - \mathsf{R})\boldsymbol{x} - \overline{\boldsymbol{z}}_{\mathsf{R}}|^p \, d\mathcal{H}^{n-1}(\boldsymbol{x}) \le c_4 \int_\Omega |\nabla \boldsymbol{\phi}(\boldsymbol{x}) - \mathsf{R}|^p \, d\boldsymbol{x},$$

hence

$$\|\mathsf{I} - \mathsf{R}\|_p^p = |\Omega| |\mathsf{I} - \mathsf{R}|^p \le c_5 \|\nabla \boldsymbol{\phi} - \mathsf{R}\|_p^p.$$

Then one has

$$\|\nabla \boldsymbol{\phi} - \mathsf{I}\|_p \le \|\nabla \boldsymbol{\phi} - \mathsf{R}\|_p + \|\mathsf{I} - \mathsf{R}\|_p \le c_6 \|\nabla \boldsymbol{\phi} - \mathsf{R}\|_p \tag{9}$$

for every $\mathsf{R} \in SO(n)$.

Now we need the celebrated *geometric rigidity lemma* by Friesecke, James & Müller [5], which holds for $1 < p < \infty$, as pointed out by Conti & Schweizer [3, Sec. 2.4]:
there is a constant $K > 0$ such that

$$\min_{\mathsf{R} \in SO(n)} \|\nabla \boldsymbol{\phi} - \mathsf{R}\|_p^p \le K \int_\Omega \operatorname{dist}^p(\nabla \boldsymbol{\phi}(\boldsymbol{x}), SO(n)) \, d\boldsymbol{x}$$

for any $\boldsymbol{\phi} \in W^{1,p}(\Omega;\mathbb{R}^n)$.

By assumption, $\det \nabla\boldsymbol{\phi} > 0$ a.e. in Ω, hence, combining Lemma 1 with the geometric rigidity lemma it follows that there exists $\mathsf{R} \in SO(n)$ such that

$$\|\nabla\boldsymbol{\phi} - \mathsf{R}\|_p^p \leq cK \int_\Omega |\nabla\boldsymbol{\phi}(\boldsymbol{x})^\mathsf{T} \nabla\boldsymbol{\phi}(\boldsymbol{x}) - \mathsf{I}|^{p/2}\, d\boldsymbol{x} = c_7 \|\nabla\boldsymbol{\phi}^\mathsf{T} \nabla\boldsymbol{\phi} - \mathsf{I}\|_{p/2}^{p/2}$$

and, by (9),

$$\|\nabla\boldsymbol{u}\|_p^p = \|\nabla\boldsymbol{\phi} - \mathsf{I}\|_p^p \leq c_6^p \|\nabla\boldsymbol{\phi} - \mathsf{R}\|_p^p \leq c_6^p c_7 \|\nabla\boldsymbol{\phi}^\mathsf{T} \nabla\boldsymbol{\phi} - \mathsf{I}\|_{p/2}^{p/2} = c\|\mathsf{E}(\boldsymbol{u})\|_{p/2}^{p/2},$$

which concludes the proof.

Remark 1 In proving the previous theorem, we also proved the following consequence of the L^p version of the geometric rigidity lemma:
there is a constant $C > 0$ *such that*

$$\|\nabla\boldsymbol{\phi} - \mathsf{I}\|_p \leq C \|\operatorname{dist}(\nabla\boldsymbol{\phi}, SO(n))\|_p$$

for every $\boldsymbol{\phi} \in W^{1,p}(\Omega;\mathbb{R}^n)$ *with* $\boldsymbol{\phi}(\boldsymbol{x}) = \boldsymbol{x}$ *on* Γ.

Acknowledgements This research is partially supported by GNFM (Gruppo Nazionale di Fisica Matematica) of INdAM (Istituto Nazionale di Alta Matematica).

References

1. Ciarlet, P.G.: Mathematical elasticity. Vol. I: Three-dimensional elasticity, *Studies in Mathematics and its Applications*, vol. 20. North-Holland Publishing Co., Amsterdam (1988)
2. Ciarlet, P.G., Mardare, C.: Continuity of a deformation in H^1 as a function of its Cauchy-Green tensor in L^1. J. Nonlinear Sci. **14**(5), 415–427 (2004). DOI 10.1007/s00332-004-0624-y.
3. Conti, S., Schweizer, B.: Rigidity and gamma convergence for solid-solid phase transitions with SO(2) invariance. Comm. Pure Appl. Math. **59**(6), 830–868 (2006). DOI 10.1002/cpa.20115.
4. Dal Maso, G., Negri, M., Percivale, D.: Linearized elasticity as Γ-limit of finite elasticity. Set-Valued Analysis **2**, 165–183 (2002).
5. Friesecke, G., James, R.D., Müller, S.: A theorem on geometric rigidity and the derivation of nonlinear plate theory from three-dimensional elasticity. Comm. Pure Appl. Math. **55**(11), 1461–1506 (2002). DOI 10.1002/cpa.10048.
6. John, F.: Bounds for deformations in terms of average strains. In: O. Shisha (ed.) Inequalities, III (Proc. Third Sympos., Univ. California, Los Angeles, Calif., 1969; dedicated to the memory of Theodore S. Motzkin), pp. 129–144. Academic Press, New York (1972)
7. Mardare, C.: A nonlinear Korn inequality with boundary conditions and its relation to the existence of minimizers in nonlinear elasticity. C. R. Math. Acad. Sci. Paris **349**(3-4), 229–232 (2011). DOI 10.1016/j.crma.2011.01.011.